## CT-BPI Competition ELISA Protocol

## **Day #1**

- 1. Coat each well with 100ul CT-BPI (10ug/ml, diluted in PBS)
- 2. Incubate for 2hr at 37C
- 3. Wash 3X with PBS + 0.05% Tween
- 4. Add 300ul PBS+5% Normal Goat Serum (NGS) and incubate o/n at RT
- 5. Incubate CF sera with peptides or w/o, in 100ul PBS + 1% NGS, at 1:100 dilution; 20ug/ml peptide o/n at 4C

## **Day #2**

- 6. Wash CT-BPI coated ELISA plate 3X with PBS + 0.05% Tween
- 7. Add 100ul patient serum that has been incubating o/n at 4C
- 8. Incubate for 1hr at RT
- 9. Wash 3X with PBS + 0.05% Tween
- 10.Add 100ul goat-anti-human HRP-labeled 2' ab (1:50,000 dil. in PBS+1% NGS) and incubate for 1hr at RT; take out Substrate Reagents and Stop soln. to warm up at RT, keep in the dark.
- 11. Wash 3X with PBS + 0.05% Tween
- 12.Add 100ul substrate (R&D, Substrate Reagent A: Substrate Reagent B 1:1)
- 13.Incubate at RT in the dark for ~15min
- 14.Add 50ul stop solution and read absorbance at 450nm and 570nm (for correction).